

# DEFAULT CLINICAL PREFERENCES

# DEFAULT CLINICAL PREFERENCES

<b>Passive aligners</b>	<ul style="list-style-type: none"><li>• Add passive aligners to match the arch with more stages</li></ul>
	<ul style="list-style-type: none"><li>• Create an equal number of active aligners on both arches</li></ul>
	<ul style="list-style-type: none"><li>• <i>The arches may have different numbers of stages</i></li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Anterior teeth leveling Upper Arch</b>	<ul style="list-style-type: none"><li>• Laterals 0.5 mm above Centrals</li><li>• Canines 0.5 mm above Laterals</li></ul>
	<ul style="list-style-type: none"><li>• Centrals and Laterals at the same level</li></ul>
	<ul style="list-style-type: none"><li>• <i>Canines and Centrals at the same level</i></li><li>• <i>Laterals 0.5 mm above</i></li></ul>
	<ul style="list-style-type: none"><li>• Follow lip guidance on frontal smiling photo</li></ul>
	<ul style="list-style-type: none"><li>• Align and level following ideal occlusion</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Expansion (teeth to move)</b>	<ul style="list-style-type: none"><li>• <i>Expand permanent teeth only</i></li></ul>
	<ul style="list-style-type: none"><li>• Expand primary and permanent teeth</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Transverse expansion</b>	<ul style="list-style-type: none"><li>• 3 to 3</li></ul>
	<ul style="list-style-type: none"><li>• 5 to 5</li></ul>
	<ul style="list-style-type: none"><li>• 7 to 7</li></ul>

*\*No default clinical preferences.*

# DEFAULT CLINICAL PREFERENCES

<b>Transverse expansion (mm)</b>	<ul style="list-style-type: none"><li>• Maintain (IPR may be needed)</li></ul>
	<ul style="list-style-type: none"><li>• <math>\leq 2</math> mm</li></ul>
	<ul style="list-style-type: none"><li>• <i>2 mm to 4 mm</i></li></ul>
	<ul style="list-style-type: none"><li>• 4 mm to 6 mm (limited)</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Curve of Spee</b>	<ul style="list-style-type: none"><li>• <i>Ideal / flat Curve of Spee (combining tipping, intrusion and extrusion)</i></li></ul>
	<ul style="list-style-type: none"><li>• Improve occlusion but do not correct the Curve of Spee unless requested on prescription.</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Class II correction</b>	<ul style="list-style-type: none"><li>• Perform sequential distalization on the upper arch</li></ul>
	<ul style="list-style-type: none"><li>• <i>Perform sequential distalization on the upper arch with IPR</i></li></ul>
	<ul style="list-style-type: none"><li>• Perform sequential mesialization on the lower arch</li></ul>
	<ul style="list-style-type: none"><li>• Perform sequential mesialization on the lower arch with IPR</li></ul>
	<ul style="list-style-type: none"><li>• Perform IPR distally to upper canines</li></ul>



# DEFAULT CLINICAL PREFERENCES

<b>Class III correction</b>	<ul style="list-style-type: none"><li>• Perform sequential mesialization on the upper arch</li></ul>
	<ul style="list-style-type: none"><li>• Perform sequential mesialization on the upper arch with IPR</li></ul>
	<ul style="list-style-type: none"><li>• Perform sequential distalization on the lower arch</li></ul>
	<ul style="list-style-type: none"><li>• <i>Perform sequential distalization on the lower arch with IPR</i></li></ul>
	<ul style="list-style-type: none"><li>• IPR distally on lower Canines</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Overjet &amp; Overbite</b>	<ul style="list-style-type: none"><li>• <i>Set to ideal overjet, with no anterior contact</i></li></ul>
	<ul style="list-style-type: none"><li>• Set overjet with slight overbite and light anterior contact</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Crowding preferences (Mild to Moderate)</b>	<ul style="list-style-type: none"><li>• <i>Expand canines and premolars</i></li><li>• <i>IPR as needed</i></li><li>• <i>Anterior protrusion</i></li></ul>
	<ul style="list-style-type: none"><li>• Expand canines, premolars and first molars</li><li>• IPR as needed</li><li>• No anterior protrusion</li></ul>

# DEFAULT CLINICAL PREFERENCES

<b>Crowding preferences (Moderate to Severe)</b>	<ul style="list-style-type: none"><li>• Expand canines and premolars, sequential distalization up to 2 to 2.5 mm (no third molars)</li><li>• Place IPR as needed, and place anterior round tripping as needed</li></ul>
	<ul style="list-style-type: none"><li>• <i>Expand canines and premolars, sequential distalization up to 2 to 2.5 mm (no third molars)</i></li><li>• <i>Place IPR as needed, but no anterior round tripping</i></li></ul>
	<ul style="list-style-type: none"><li>• Expand canines and premolars, sequential distalization up to 2 to 2.5 mm (no third molars)</li><li>• Place anterior round tripping as needed, but no IPR</li></ul>

# DEFAULT CLINICAL PREFERENCES

IPR on primary teeth	Yes
	<i>No</i>

# DEFAULT CLINICAL PREFERENCES

<b>Start IPR</b>	<ul style="list-style-type: none"><li>• Fully align teeth before IPR</li></ul>
	<ul style="list-style-type: none"><li>• Start IPR at the beginning of the treatment</li></ul>
	<ul style="list-style-type: none"><li>• Start IPR from step 2</li></ul>
	<ul style="list-style-type: none"><li>• Start IPR from step 3</li></ul>
	<ul style="list-style-type: none"><li>• <i>IPR whenever it's necessary</i></li></ul>
	<ul style="list-style-type: none"><li>• I prefer to avoid IPR</li></ul>

# DEFAULT CLINICAL PREFERENCES

IPR Maximum	<ul style="list-style-type: none"><li>• 0.2 mm for anterior teeth / 0.5 mm for posterior teeth</li></ul>
	<ul style="list-style-type: none"><li>• <i>0.4 mm for anterior teeth / 0.5 mm for posterior teeth</i></li></ul>
	<ul style="list-style-type: none"><li>• 0.5 mm for all teeth</li></ul>